

BINARY TREES AND FIBRED CATEGORIES

N. RAGHAVENDRA

ABSTRACT. We develop a purely set-theoretic formalism for binary trees and binary graphs. We define a category of binary automata, and display it as a fibred category over the category of binary graphs. We also relate the notion of binary graphs to transition systems, which arise in the theory of concurrent computing.

CONTENTS

1. Introduction	1
2. Binary graphs	2
3. Bit strings	3
4. Action of bit strings	4
5. Transition diagrams	6
6. Binary forests and trees	7
7. Binary automata and transition systems	10
8. Fibred categories	11
9. A criterion for fibredness	13
10. Split categories	14
11. Fibred category of binary automata	17
12. Conclusion	19
Acknowledgements	19
References	19

1. INTRODUCTION

This work arose out of certain issues which we encountered while trying to understand certain well-known algorithms in computational biology, whose aim is to build trees out of various data. We will discuss these issues presently. In any case, they motivated us to develop an abstract formalism for binary trees, which is the subject of this article. We found that our formalism fits neatly into the framework of fibred categories, which were originally studied in algebraic geometry. Moreover, it has connections with the notion of transition systems which arise in the theory of concurrent computation.

The problems which motivated this work are as follows. First, there are several statements in tree-construction algorithms which assert the uniqueness of some tree. One can ask certain natural questions in such a situation. To begin with, what does one mean by uniqueness: for instance, does one mean uniqueness in the sense of equality of trees, or does one mean uniqueness up to an isomorphism? Next, what is the precise property of trees, which makes the tree in question unique? Can one formulate this property as a universal property in a suitable category? Thus, we were led to seek an abstract framework for binary trees.

Secondly, the standard descriptions of trees are either recursive, as in [Knu97, Section 2.3], or are based on graph theory, as in [AHU74, Section 2.4]. We feel that these descriptions have certain

2000 *Mathematics Subject Classification*. Primary 05C05; Secondary 18D30, 05C62, 18B20, 68Q85.

Key words and phrases. Binary tree, fibred category, binary graph, binary automaton, transition system.

drawbacks. On the one hand, a recursive definition of trees sets them apart from other important classes of mathematical objects such as groups, finite automata etc., which are described in terms of sets and maps between them, whereas a set-theoretic description of trees would bring them into a general framework, consistent with the definitions of several other mathematical structures. On the other hand, a treatment of trees based on graph theory entails carrying, to some extent, the formalism of the latter subject, which can be avoided if we, *ab initio*, develop a definition of trees which does not rely on graph theory.

In view of the above discussion, in this article we develop a purely set-theoretic treatment of binary trees. We will also discuss the relationship between our definitions, and the standard graph-theoretic notions regarding trees.

We now sketch the contents of the various sections. In Section 2, we develop a set-theoretic formalism for binary graphs. In Section 3, we establish our notation for the free monoid on the set of bits, which monoid acts on the set of nodes of any binary graph, as described in Section 4. Transition diagrams, defined in Section 5, establish a connection between our model for binary graphs and their standard graph-theoretic definition. In Section 6, we describe binary trees as a specialization of binary forests. We introduce the category of binary automata in Section 7, and we relate binary graphs to transition systems. The important notion of a fibred category is recalled in Section 8, and in the two subsequent Sections, we recall a criterion for a category to be fibred, and the notion of split categories. To assist the reader, and to make this paper self-contained, we give complete definitions and proofs regarding fibred categories, even though the material may be well-known to specialists in certain areas of algebraic geometry or category theory. In Section 11, we display the category of binary automata as a fibred category over the category of binary graphs. In Section 12, we summarise the article, and mention some ideas for future work.

2. BINARY GRAPHS

We begin our discussion of binary trees with the definition of a more general object, which we call a *binary graph*. This helps us to introduce several constructions which are useful in our later discussion of binary trees.

Definition 2.1. A *binary graph* is a pair $\Gamma = (Q, \delta)$, where

- (1) Q is a finite set, whose elements are called *nodes*.
- (2) $\delta : Q \times \Sigma \rightarrow Q$ is a function, called the *transition function*, where $\Sigma = \{0, 1\}$.

Given such a binary graph Γ , we get a pair of functions $\lambda : Q \rightarrow Q$ and $\rho : Q \rightarrow Q$, defined by

$$\lambda(q) = \delta(q, 0), \quad \text{and} \quad \rho(q) = \delta(q, 1), \quad \text{for all } q \in Q.$$

We call $\lambda(q)$ the *left child* of q , and $\rho(q)$ the *right child* of q .

Definition 2.2. Let $\Gamma = (Q, \delta)$ and $\Gamma' = (Q', \delta')$ be binary graphs. A *morphism* from Γ to Γ' is a function $f : Q \rightarrow Q'$, such that $f(\delta(q, a)) = \delta'(f(q), a)$, for all $q \in Q$ and for all letters $a \in \Sigma$. We say that Γ is a *binary subgraph* of Γ' if Q is a subset of Q' , and if the inclusion $i : Q \hookrightarrow Q'$ is a morphism of binary graphs.

Example 2.3. Let $Q = \{q_0, q_1\}$, and define $\delta : Q \times \Sigma \rightarrow Q$ by $\delta(q, 0) = q_0$ and $\delta(q, 1) = q_1$, for all $q \in Q$. It is convenient to define the transition function δ through a table, whose rows are indexed by the letters in Σ , whose columns are indexed by the nodes in Q , and whose (a, q) -th entry equals $\delta(q, a)$. In this notation, the above function is defined by the following table:

	q_0	q_1
0	q_0	q_0
1	q_1	q_1

Example 2.4. Let $Q = \{q_0, q_1, q_2, q_3, q_4, q_5, q_6\}$, and define $\delta : Q \times \Sigma \rightarrow Q$ by the following table:

	q_0	q_1	q_2	q_3	q_4	q_5	q_6
0	q_1	q_3	q_4	q_3	q_4	q_6	q_6
1	q_2	q_4	q_5	q_3	q_4	q_5	q_6

Example 2.5. Let $Q = \{q_0, q_1, q_2, q_3, q_4, q_5, q_6, q_7\}$, and define $\delta : Q \times \Sigma \rightarrow Q$ by the following table:

	q_0	q_1	q_2	q_3	q_4	q_5	q_6	q_7
0	q_1	q_3	q_2	q_3	q_4	q_6	q_6	q_7
1	q_2	q_4	q_2	q_3	q_4	q_7	q_6	q_7

Example 2.6. Let $Q = \{q_0, q_1, q_2, q_3\}$, and define $\delta : Q \times \Sigma \rightarrow Q$ by the following table:

	q_0	q_1	q_2	q_3
0	q_1	q_3	q_2	q_3
1	q_2	q_1	q_2	q_3

3. BIT STRINGS

In this section, we establish our notation for the languages on the set of bits. Let Σ^* be the free monoid on the set Σ which, we recall, is the set $\{0, 1\}$. Thus, Σ^* is a disjoint union,

$$\Sigma^* = \coprod_{n=0}^{\infty} \Sigma^n,$$

where Σ^0 is a singleton consisting of a symbol ϵ , and $\Sigma^n = \Sigma \times \cdots \times \Sigma$ (n times). We call elements of Σ^* *words* or *bit strings*. In particular, the word ϵ is called the *empty word*. We call the set Σ an *alphabet*, and refer to the elements of Σ as *letters*. If a word $w \in \Sigma^n$, we say that w has length n , and write $|w| = n$. Following standard practice, we denote a word $w = (a_1, \dots, a_n) \in \Sigma^n$ by $w = a_1 \cdots a_n$. There is a natural multiplication law in the set Σ^* ,

$$\begin{aligned} \Sigma^* \times \Sigma^* &\rightarrow \Sigma^*, \\ (v, w) &\mapsto vw, \end{aligned}$$

which is given by concatenation of words. Explicitly, we define

$$\epsilon w = w \epsilon = w \quad \text{for all } w \in \Sigma^*,$$

and if $v = a_1 \cdots a_m$ and $w = b_1 \cdots b_n$, where $a_i, b_j \in \Sigma$ for all $i = 1, \dots, m$ and $j = 1, \dots, n$, then we set

$$vw = a_1 \cdots a_m b_1 \cdots b_n.$$

With this multiplication law, the set Σ^* becomes a monoid, whose identity element is the empty word ϵ . It is clear that $|vw| = |v| + |w|$ for all $v, w \in \Sigma^*$. The monoid Σ^* is free on the set Σ , i.e., it has the following universal property: given any monoid M , and given any function $f : \Sigma \rightarrow M$, there exists a unique homomorphism of monoids, $\tilde{f} : \Sigma^* \rightarrow M$, which extends f , i.e., which makes the diagram

$$\begin{array}{ccc} \Sigma & \xrightarrow{i} & \Sigma^* \\ & \searrow f & \downarrow \tilde{f} \\ & & M \end{array}$$

commute, where $i : \Sigma \rightarrow \Sigma^*$ is the natural inclusion map.

4. ACTION OF BIT STRINGS

Let $\Gamma = (Q, \delta)$ be a binary graph. The universal property of Σ^* implies that there is a unique right action of the monoid Σ^* on Q ,

$$\begin{aligned}\hat{\delta} : Q \times \Sigma^* &\rightarrow Q, \\ (q, w) &\mapsto qw,\end{aligned}$$

such that

$$\hat{\delta}(q, aw) = \hat{\delta}(\delta(q, a), w) \quad \text{for all } q \in Q, a \in \Sigma, \text{ and } w \in \Sigma^*.$$

By abuse of notation, we denote $\hat{\delta}$ also by δ .

For any node $q \in Q$, let

$$\Sigma_q^* = \{w \in \Sigma^* \mid qw = q\}$$

be the isotropy submonoid of Σ^* at q . We say that a node q is a *leaf* if it is fixed by Σ^* , i.e., if $\Sigma_q^* = \Sigma^*$. If q is not a leaf, i.e., if Σ_q^* is a proper submonoid of Σ^* , then we call q an *internal node* of the binary graph Γ . Note that q is a leaf if and only if $\delta(q, a) = q$ for all letters $a \in \Sigma$.

Example 4.1. The binary graph of Example 2.3 has no leaves. In Example 2.4, the nodes q_3, q_4 and q_6 are the leaves. The leaves of Γ , in Example 2.5, are the nodes q_2, q_3, q_4, q_6 and q_7 .

We say that an element q of Q is a *parent* of q' if $q \neq q'$, and $\delta(q, a) = q'$ for some $a \in \Sigma$. In this case q has to be an internal node. A *path* in Γ is a pair $P = (q_0, w)$, where $q_0 \in Q$ is a node, and $w \in \Sigma^*$ is a word, satisfying one of the following two conditions:

- (1) Either $w = \epsilon$ is the empty word. Or,
- (2) $w = a_1 \cdots a_n$, with $a_i \in \Sigma$, and $q_i \neq q_{i-1}$, where $q_i = q_0 a_1 \cdots a_i$, for each $i = 1, \dots, n$. (In particular, q_{i-1} is a parent of q_i , for each $i = 1, \dots, n$.)

We say that this path *starts* at q_0 , *ends* at $q_0 w$, and has *length* $|w|$. A *trivial path* is one which has length 0; it has to be of the form (q_0, ϵ) , for some node q_0 . We say that q is an *ancestor* of q' , or that q' is a *descendant* of q , if there is a path starting at q and ending at q' . If, in addition, $q \neq q'$, then we say that q is a *proper ancestor* of q' , or that q' is a *proper descendant* of q .

Definition 4.2. Let $\Gamma = (Q, \delta)$ be a binary graph. For any pair of nodes, $q, q' \in Q$, we write $q' \leq q$ (respectively, $q' < q$) if q is an ancestor (respectively, a proper ancestor) of q' . We write $q \geq q'$ (respectively, $q > q'$) if $q' \leq q$ (respectively, $q' < q$). The relation $\leq$ on Q is a preorder, i.e., it has all the properties of a partial order, except that of anti-symmetry. In other words, the relation $\leq$ is reflexive and transitive, but there may exist elements $q, q' \in Q$ such that $q \leq q'$ and $q' \leq q$, and yet $q \neq q'$. We call this preorder the *inheritance preorder* on Q .

Remark 4.3. Any path which starts at a leaf is a trivial path.

Remark 4.4. Let $\Gamma = (Q, \delta)$ and $\Gamma' = (Q', \delta')$ be binary graphs. A morphism from Γ to Γ' , introduced in Definition 2.2, is just a function $f : Q \rightarrow Q'$, which is Σ^* -equivariant, i.e., which satisfies the condition that $f(qw) = f(q)w$ for all $q \in Q$ and $w \in \Sigma^*$. Similarly, if $\Gamma' = (Q', \delta')$ is a binary graph, and Q is a Σ^* -invariant subset of Q' , which means that $qw \in Q$ for all $q \in Q$ and for all $w \in \Sigma^*$, we obtain a binary subgraph $\Gamma = (Q, \delta)$ of Γ' , where $\delta : Q \times \Sigma \rightarrow Q$ is the restriction of $\delta' : Q' \times \Sigma \rightarrow Q'$. Thus, binary subgraphs of Γ' are in bijective correspondence with Σ^* -invariant subsets of Q' . Note that a subset Q of Q' is Σ^* -invariant if and only if $\delta'(q, a) \in Q$, for all $q \in Q$ and for all letters $a \in \Sigma$.

For every node $q \in Q$, let

$$\Sigma^*(q) = \{qw \mid w \in \Sigma^*\}$$

be the orbit of q .

Proposition 4.5. In any binary graph $\Gamma = (Q, \delta)$, a node q is an ancestor of q' if and only if $q' \in \Sigma^*(q)$.

Before proving the Proposition, let us introduce a device which will be useful in the proof. Let $\Gamma = (Q, \delta)$ be a binary graph, and let $w \in \Sigma^*$. For each node $q_0 \in Q$, we shall define a new word $w[q_0]$ as follows. If $w = \epsilon$, then set $w[q_0] = w = \epsilon$. If w is nonempty, write $w = a_1 \cdots a_n$, where $a_i \in \Sigma$, and let $q_i = q_0 a_1 \cdots a_i$, for $i = 1, \dots, n$. If $q_i = q_{i-1}$ for all $i = 1, \dots, n$, define $w[q_0] = \epsilon$. If not all q_i are equal, let $i_1 < \cdots < i_k$ be the elements of the set

$$\{i \mid 1 \leq i \leq n \text{ and } q_i \neq q_{i-1}\}.$$

Let $b_r = a_{i_r}$, for $r = 1, \dots, k$, and define $w[q_0] = b_1 \cdots b_k$. The words w , $w[q_0]$ and the indices i_r have the following properties:

- (1) The pair $P[q_0, w] = (q_0, w[q_0])$ is a path, starting at q_0 and ending at $q_0 w$.
- (2) We have

$$\begin{aligned} q_i &= q_0, & \text{if } 0 \leq i < i_1, \\ q_i &= q_{i_r}, & \text{if } 1 \leq r < k \text{ and } i_r \leq i < i_{r+1}, \text{ and} \\ q_i &= q_{i_k}, & \text{if } i_k \leq i \leq n. \end{aligned}$$

- (3) The letters a_i satisfy the following conditions:

$$\begin{aligned} q_0 a_i &= q_0, & \text{if } 1 \leq i < i_1, \\ q_{i_r} a_i &= q_{i_r}, & \text{if } 1 \leq r < k \text{ and } i_r < i < i_{r+1}, \text{ and} \\ q_{i_k} a_i &= q_{i_k}, & \text{if } i_k < i \leq n. \end{aligned}$$

- (4) The b_i satisfy the following equations:

$$\begin{aligned} q_0 b_1 &= q_{i_1}, \\ q_{i_r} b_{r+1} &= q_{i_{r+1}}, & \text{if } 1 \leq r < k. \end{aligned}$$

Definition 4.6. Let $\Gamma = (Q, \delta)$ be a binary graph, and let $w \in \Sigma^*$ be a word. Then, for each node $q_0 \in Q$, we call the word $w[q_0]$ defined above, the *contraction* of w by q_0 .

Proof of Proposition 4.5. Clearly, if there is a path from q to q' , then $q' \in \Sigma^*(q)$. Conversely, suppose that $q' \in \Sigma^*(q)$, say $q' = qw$. Then, by the device introduced above, there exists a word $w[q]$ such that the pair $P = (q, w[q])$ is a path from q to $qw = q'$. $\square$

Remark 4.7. If q is a leaf in a binary graph $\Gamma = (Q, \delta)$, then Σ^* fixes q , so we get a binary subgraph $\Gamma_q = (\{q\}, \delta_q)$, where δ_q is the unique map $\{q\} \times \Sigma \rightarrow \{q\}$. More generally, if q is any node, then we obtain a binary subgraph $\Gamma_q = (\Sigma^*(q), \delta_q)$, where $\Sigma^*(q)$ is the orbit of q , and $\delta_q : \Sigma^*(q) \times \Sigma \rightarrow \Sigma^*(q)$ is the restriction of $\delta : Q \times \Sigma \rightarrow Q$. We call Γ_q the *orbit graph* of q . It is obvious that Γ_q is a minimal binary subgraph of Γ , whenever q is a leaf of Γ .

The next proposition provides criteria, in terms of the action of Σ^* , for the anti-symmetry of the inheritance preorder $\leq$, defined above (see Definition 4.2). We say that a word $v \in \Sigma^*$ is a *prefix* of another word w , if $w = vz$ for some word $z \in \Sigma^*$. In this case we say that the word z is a *suffix* of w . We say that v is a *proper prefix* (respectively, *proper suffix*) of w if v is a prefix (respectively, suffix) of w , $v \neq \epsilon$, and $v \neq w$. A subset L of Σ^* is said to be *prefix-closed*, if whenever $w \in L$, every prefix v of w also belongs to L . The notion of a *suffix-closed* subset of Σ^* is defined in a similar manner. Clearly, every nonempty prefix-closed set has to contain the empty word ϵ , and the singleton $\{\epsilon\}$ is the smallest nonempty prefix-closed subset of Σ^* . An analogous observation applies to suffix-closed sets, too.

We say that a subset L of Σ^* is *subword-closed* if, whenever $w_1, w_2, \dots, w_n$ are elements of Σ^* , whose product $w_1 w_2 \cdots w_n$ belongs to L , we have $w_i \in L$ for all $i = 1, \dots, n$. This is equivalent to the condition that L is both prefix-closed and suffix-closed.

A *cycle* in a binary graph $\Gamma = (q, \delta)$ is a path $P = (q_0, w)$, such that $q_0 w = q_0$. We say that Γ is *acyclic* if every cycle in Γ is a trivial path.

Proposition 4.8. *Let $\Gamma = (Q, \delta)$ be a binary graph. Then, the following statements are equivalent:*

- (1) *The inheritance preorder $\leq$ on Q (see Definition 4.2) is a partial order.*
- (2) *For every node $q \in Q$, the isotropy submonoid Σ_q^* is prefix-closed.*
- (3) *For every node $q \in Q$, the isotropy submonoid Σ_q^* is subword-closed.*
- (4) *For every internal node $q \in Q$, the isotropy submonoid Σ_q^* is one of the three submonoids $\{\epsilon\}$, $\{0\}^*$ and $\{1\}^*$.*
- (5) *The binary graph Γ is acyclic.*

Proof. Suppose, first, that the inheritance preorder, $\leq$, is anti-symmetric. Let $q \in Q$ be a node, let $w \in \Sigma_q^*$, and let v be a prefix of w , say $w = vz$, for some word z . Then the node $q' = qv$ clearly satisfies the inequality $q' \leq q$. On the other hand, $q = q'z$, so $q \leq q'$. Therefore, by the anti-symmetry of $\leq$, we obtain $q' = q$. Since $q = q' = qv$, the word v belongs to Σ_q^* and, hence, that set is prefix-closed. We have, thus, proved that the condition (1) implies (2).

Let us, now, suppose that Σ_q^* is prefix-closed, and that $w_1, \dots, w_n$ are words, whose product $w = w_1 \cdots w_n$ belongs to Σ_q^* . Since the isotropy submonoid is prefix-closed, it contains, for each $i = 1, \dots, n$, the prefixes $w_1 \cdots w_{i-1}$ and $w_1 \cdots w_i$, of w . Therefore,

$$qw_i = (qw_1 \cdots w_{i-1})w_i = q(w_1 \cdots w_{i-1}w_i) = q,$$

i.e., $w_i \in \Sigma_q^*$, for all $i = 1, \dots, n$. So, Σ_q^* is subword-closed. Conversely, every subword-closed set is, obviously, prefix-closed. This proves that (2) and (3) are equivalent statements.

The conditions (3) and (4) are equivalent, for the only subword-closed proper submonoids of Σ^* are $\{\epsilon\}$, $\{0\}^*$ and $\{1\}^*$.

Assume that Γ satisfies the condition (2), and let $P = (q_0, w)$ be a cycle in Γ . Suppose that P has positive length n , say, $w = a_1 \cdots a_n$, where $a_i \in \Sigma$. Let $q_i = q_0 a_1 \cdots a_n$, for $i = 1, \dots, n$. Since P is a cycle, $q_0 w = q_0$, hence $w \in \Sigma_{q_0}^*$. The set $\Sigma_{q_0}^*$ is prefix-closed, so it contains the prefix a_1 of w . Therefore, $q_1 = q_0 a_1 = q_0$, a contradiction since q_{i-1} is a parent of q_i for all i . The cycle P , therefore, has to be a trivial path, so (2) implies (5).

Suppose, finally, that Γ is an acyclic binary graph, and let $q, q' \in Q$ be such that $q \leq q'$ and $q' \leq q$. Since $q' \leq q$, by the definition of the partial order on Q , there exists a path $P = (q_0, w)$ in Γ such that $q_0 = q$, and $q_0 w = q'$. Since $q \leq q'$, there exists a path $P' = (q'_0, w')$ such that $q'_0 = q'$ and $q'_0 w' = q$. The pair PP' , defined by $PP' = (q_0, ww')$, is a path, and is, in fact, a cycle in Γ . Since Γ is acyclic, the cycle PP' must have length 0, hence $w = w' = \epsilon$. Therefore, $q' = q_0 w = q_0 = q$ and, so, the preorder $\leq$ is anti-symmetric. We have, now, shown that the condition (5) implies (1). $\square$


5. TRANSITION DIAGRAMS

Transition diagrams are a device through which we connect our description of binary graphs with their natural treatment in graph theory. Let $\Gamma = (Q, \delta)$ be a binary graph. Then the *transition diagram* of Γ is a labelled directed graph $D(\Gamma)$, which is defined as follows:


- (1) The vertex set of $D(\Gamma)$ is defined to be Q .
- (2) If $(q, q', a) \in Q \times Q \times \Sigma$, if $q \neq q'$, and if $\delta(q, a) = q'$, then there is an edge labelled a from q to q' .

Thus, there is an edge from q to q' if and only if q is a parent of q' . If $(q, a) \in Q \times \Sigma$, and if there is no edge from q labelled a , then it means that $\delta(q, a) = q$. In particular, the vertices of $D(\Gamma)$ which have no edge starting from them are precisely the leaves in Q .


Example 5.1. The following directed graph is the transition diagram of the binary graph of Example 2.3.




Example 5.2. The transition diagram of the binary graph defined in Example 2.4, is as follows:



Example 5.3. The binary graph of Example 2.5 has the following transition diagram:



Example 5.4. Here is the transition diagram of the binary graph described in Example 2.6:



6. BINARY FORESTS AND TREES

In this Section, we will define binary forests and trees, in our formalism, and discuss their basic properties in terms of the inheritance preorder and the action of bit strings.

Definition 6.1. A *binary forest* is a binary graph $\Gamma = (Q, \delta)$, which satisfies the following conditions:

- (1) The inheritance preorder on Q is a partial order (see Definition 4.2 and Proposition 4.8).
- (2) If $qa = q'b$, where a and b are letters in Σ , then either $qa = q$, or $q'b = q$, or $(q, a) = (q', b)$.

Example 6.2. The condition (2) in Definition 6.1 can be restated as the condition that every node can have at most one parent. For instance, the binary graph in Example 2.5 satisfies this condition, and is a binary forest. The graph of Example 2.4 does not satisfy the condition (2). The graph in Example 2.3 does not satisfy the condition (1). Note that every subgraph of a binary forest is, again, a binary forest. We will sometimes refer to such a subgraph as a *subforest*.

We say that a path $P = (q_0, w)$ is a *suffix* of a path $P' = (q'_0, w')$, if there exists a word $v \in \Sigma^*$, such that $w' = vw$, and $q_0 = q'_0v$. Note that if this is the case, then P and P' end at the same node.

Proposition 6.3. A binary graph $\Gamma = (Q, \delta)$ satisfies the condition (2) of Definition 6.1, if and only if it satisfies the following condition:

(2') If P and P' are paths in Γ , which end at the same node, then either P is a suffix of P' , or P' is a suffix of P .

Proof. Suppose that Γ satisfies the condition (2') of the Proposition. Let q, a, q', b be as in (2) of Definition 6.1, and suppose that $qa \neq q$ and $q'b \neq q'$. Then $P = (q, a)$ and $P' = (q', b)$ are paths in Γ which end at the same node. Therefore, by (2'), one of them is a suffix of the other. As the both have the same length, 1, this means that they are equal, i.e., $q = q'$ and $a = b$, so, (2') implies condition (2) of Definition 6.1.

Conversely, assume that Γ satisfies the condition (2) of Definition 6.1. Let $P = (q_0, w)$ and $P' = (q'_0, w')$ be two paths, which end at the same node, i.e., $q_0w = q'_0w'$. It is obvious that if w is the empty word, then P is a suffix of P' and, similarly, if w' is the empty word, then P' is a suffix of P . So, let us assume that both w and w' have positive length, say, $w = a_1 \cdots a_m$ and $w' = a'_1 \cdots a'_n$. Let $q_i = q_0a_1 \cdots a_i$ and $q'_j = q'_0a'_1 \cdots a'_j$, for $i = 1, \dots, m$ and $j = 1, \dots, n$. Suppose, without loss of generality, that $m \leq n$. Since P and P' are paths, we have $q_{m-1} \neq q_m$ and $q'_{n-1} \neq q'_n$. So, applying the condition (2) to the equation $q_{m-1}a_m = q'_{n-1}a'_n$, we get that $q_{m-1} = q'_{n-1}$ and $a_m = a'_n$. Considering, now, the paths $(q_0, a_1 \cdots a_{m-1})$ and $(q'_0, a'_1 \cdots a'_{n-1})$, and using induction, we see that $q_i = q'_{n-m+i}$ for all $i = 0, 1, \dots, m$, and $a_i = a'_{n-m+i}$ for all $i = 1, \dots, m$. Setting $v = a'_1 \cdots a'_{n-m}$, we have $w' = vw$, and $q'_0v = q'_{n-m} = q_0$. Therefore, P is a suffix of P' . $\square$

Corollary 6.4. Let $\Gamma = (Q, \delta)$ be a binary forest. Then, for every pair of nodes $q, q' \in Q$ such that $q' \leq q$, there is a unique path from q to q' .

Proof. Let $P = (q, w)$ and $P' = (q, w')$ be two paths from q to q' . Then by Proposition 6.3, one of these two paths is a suffix of the other, say, P is a suffix of P' . Thus, there exists a word $v \in \Sigma^*$, such that $w' = vw$, and $q = qv$. Since P' is a path, so is the pair $Q = (q, v)$, and clearly Q starts and ends at q . By Proposition 4.8, Γ is acyclic, hence the cycle Q is a trivial path. Therefore, $v = \epsilon$, which implies that $w' = w$ and $P' = P$. $\square$

Let X be a set with a preorder $\leq$, and let S be a subset of X . Recall that an element x_0 of S is said to be a *maximum* element of S if $x \leq x_0$ for all $x \in S$, and that x_0 is said to be a *maximal* element of S if, whenever $x \in S$ and $x_0 \leq x$, we have $x = x_0$.

Remark 6.5. Note that if the preorder $\leq$ on X is a partial order, then a subset of X has at most one maximum. Also, in any preorder, if a set has a maximal element and a maximum element, then they must coincide, and the set has a unique maximum.

Proposition 6.6. Consider a binary graph $\Gamma = (Q, \delta)$, and the inheritance preorder, $\leq$, on Q (see Definition 4.2). Then, the following statements are true:

- (1) The set of nodes, Q , has a maximal element, and is the union of the Σ^* -orbits of all the maximal elements.
- (2) If Γ is a binary forest, and if q_1 and q_2 are distinct maximal elements of Q , then their orbits, $\Sigma^*(q_1)$ and $\Sigma^*(q_2)$ are disjoint.

Proof. Let $q_0 \in Q$ be a node. For each $i \in \mathbb{N}$, inductively define a node q_i , as follows: $q_i = q_{i-1}$ if q_{i-1} is maximal, and if q_{i-1} is not maximal, then q_i is some choice of a node q such that $q_{i-1} < q$. Since Q is a finite set, the sequence $\{q_i\}_{i=0}^\infty$ is eventually constant, i.e., there exists $n \geq 0$ such that $q_i = q_n$ for all $i \geq n$. Then, q_n is a maximal node, and $q_0 \leq q_n$, i.e., $q_0 \in \Sigma^*(q_n)$. This proves that (1) is true for any binary graph.

Suppose, next, that q_1 and q_2 are maximal elements in a binary forest Γ , and that $q \in \Sigma^*(q_1) \cap \Sigma^*(q_2)$. Then, there exist a path P_1 from q_1 to q , and a path P_2 from q_2 to q . Since P_1 and P_2 end at the same node, one of them is a suffix of the other, say, P_1 is a suffix of P_2 . It follows that $q_1 \in \Sigma^*(q_2)$, so, $q_1 \leq q_2$. By maximality of q_1 , we obtain that $q_1 = q_2$. Therefore, the orbits of two distinct maximal elements in Γ cannot intersect. $\square$

Definition 6.7. Let $\Gamma = (Q, \delta)$ be a binary graph. We say that a subset S of Q is a *connected set* if it has a maximum element. A *connected component* of Γ is a maximal connected subset of Q . We say that Γ is *connected* if its set of nodes, Q , is a connected set.

If $\Gamma = (Q, \delta)$ is a connected binary graph, then Q has a unique maximum. For, by (1) of Proposition 6.6, Q has a maximal element. By Remark 6.5, this element has to be the unique maximum of Q . We call it the *root* of Γ .

Suppose that S is a connected component of a binary graph $\Gamma = (Q, \delta)$, and let q_0 be a maximum element of S . By (1) of Proposition 6.6, there exists a maximal node q_1 in Q , such that $q_0 \leq q_1$. For every element $q \in S$, we have $q \leq q_0 \leq q_1$, hence S is contained in the orbit $\Sigma^*(q_1)$. This orbit is a connected set, having q_1 as a maximum. Therefore, by the maximality of S , we have $S = \Sigma^*(q_1)$. Since S has a maximum element q_0 and a maximal element q_1 , by Remark 6.5, we have $q_0 = q_1$, and q_0 is the unique maximum of S . In other words, every connected component S of a binary graph has a unique maximum q_0 , and this maximum is a maximal element in the full set of nodes; moreover, $S = \Sigma^*(q_0)$. Therefore, by Remark 4.7, we have a binary subgraph $\Gamma_{q_0} = (\Sigma^*(q_0), \delta_{q_0})$ of Γ , having $S = \Sigma^*(q_0)$ as its set of nodes. We denote the graph Γ_{q_0} by Γ_S and, by abuse of language, we refer to Γ_S also as a connected component of Γ .

Proposition 6.8. Let $\Gamma = (Q, \delta)$ be a binary graph. Then, the orbit of any maximal node $q_0 \in Q$ is a connected component of Γ . Conversely, every connected component S of Γ has a unique maximum q_0 ; this maximum is a maximal element in the full set of nodes, and S is the orbit of q_0 .

Proof. Suppose that q_0 is a maximal element of Q . Obviously, q_0 is a maximum element of $\Sigma^*(q_0)$, so the orbit of q_0 is a connected set. Let S be a connected subset of Q , which contains $\Sigma^*(q_0)$, and denote by q_1 a maximum element of S . Then, $q_0 \leq q_1$, so, by maximality of q_0 , we have $q_0 = q_1$. Since q_0 is a maximum of S , we have $q \leq q_0$ for all $q \in S$, i.e., $q \in \Sigma^*(q_0)$ for all $q \in S$. Therefore, $S = \Sigma^*(q_0)$, hence, $\Sigma^*(q_0)$ is a maximal connected subset of Q . This proves the first assertion of the Proposition. We have already shown, above, the truth of the second assertion. $\square$

Remark 6.9. If $\Gamma = (Q, \delta)$ is a binary graph, then, for any node $q_0 \in Q$, we define

$$\begin{aligned} Q_\lambda(q_0) &= \{q_0 0v \mid v \in \Sigma^*\}, \quad \text{and} \\ Q_\rho(q_0) &= \{q_0 1v \mid v \in \Sigma^*\}. \end{aligned}$$

Since these two sets are Σ^* -invariant, they define, by Remark 4.4, two subgraphs $\Gamma_\lambda(q_0) = (Q_\lambda(q_0), \delta)$ and $\Gamma_\rho(q_0) = (Q_\rho(q_0), \delta)$, where, by abuse of notation, we denote the restriction of δ to a subset of $Q \times \Sigma$ by the same symbol. It is clear that $\delta(q_0, 0)$ is a maximum element of $Q_\lambda(q_0)$, so $\Gamma_\lambda(q_0)$ is a connected graph. We call it the *left subgraph* at q_0 . Similarly, $\Gamma_\rho(q_0)$ is also a connected graph, having $\delta(q_0, 1)$ as the maximum element. We call it the *right subgraph* at q_0 .

Definition 6.10. A connected binary forest is called a *binary tree*. In this case, the set Q has a unique maximum (see Remark 6.5), and this element is called the *root* of Γ .

Example 6.11. The binary graph in Example 2.5 is not connected, because the subset $\{q_0, q_5\}$ of Q does not have an upper bound. In this graph, q_0 and q_5 are maximal nodes. In Example 2.3, there are no maximal nodes, but both the nodes q_0 and q_1 are maximum nodes. The graph in Example 2.6 is a binary tree, with root q_0 .

Let $\Gamma = (Q, \delta)$ is a binary forest. Then, every connected component of Γ is a binary tree. As we had observed in Example 6.2, every subgraph of Γ is also a binary forest. In particular, for any node $q_0 \in Q$, the left and right subgraphs at q_0 (See Remark 6.9), namely $\Gamma_\lambda(q_0)$ and $\Gamma_\rho(q_0)$, are binary forests. Since they are, always, connected graphs, it follows that $\Gamma_\lambda(q_0)$ and $\Gamma_\rho(q_0)$ are, in fact, binary trees. We call them, respectively, the *left subtree* and the *right subtree* at q_0 .

Proposition 6.12. Let $\Gamma = (Q, \delta)$ be a binary forest, and suppose that q_0 is a node in Q which has trivial isotropy. Then, the left and right subtrees of q_0 are disjoint, i.e., $Q_\lambda(q_0) \cap Q_\rho(q_0) = \emptyset$.

Proof. Suppose that $Q_\lambda(q_0) \cap Q_\rho(q_0) \neq \emptyset$, say $q_0 0 v = q_0 1 w$ for some words $v, w \in \Sigma^*$. Let $q' = q_0 0 v$. Then, q' is a descendant of both $q_0 0$ and $q_0 1$. Therefore, by Proposition 4.5, there exist paths $P_0 = (q_0 0, z_0)$ and $P_1 = (q_0 1, z_1)$ from $q_0 0$ and $q_0 1$, respectively, to q' . Since q_0 has trivial isotropy, $q_0 0 \neq q_0$ and $q_0 1 \neq q_0$. Therefore, $Q_0 = (q_0, 0z_0)$ and $Q_1 = (q_0, 1z_1)$ are also paths. Since both of them start at the same point q_0 , and end at the same point q' , by Corollary 6.4, $Q_0 = Q_1$, i.e., $0z_0 = 1z_1$. Since this is an impossibility, we conclude that our hypothesis, that the left and right subtrees of q_0 overlap, has to be false. $\square$

7. BINARY AUTOMATA AND TRANSITION SYSTEMS

We will, now, recall the notion of binary automata, and describe a canonical functor from the category of these automata, to that of binary graphs with a distinguished node. We will also relate binary graphs to transition systems.

A *binary automaton* is a finite automaton $M = (Q, \Sigma, \delta, q_0, F)$, where the input alphabet Σ equals the set $\{0, 1\}$. (We follow the notation of [HU79, Chapter 1] for finite automata.) Let us define a state q to be *stationary* if $\delta(q, a) = q$ for all $a \in \Sigma$. We will say that a binary automaton is *admissible*, in case the set F of final states coincides with the set of stationary states.

Let $M = (Q, \Sigma, \delta, q_0, F)$ and $M' = (Q', \Sigma, \delta', q'_0, F')$ be binary automata. A *morphism* from M to M' is a function $f : Q \rightarrow Q'$, such that the following conditions hold:

- (1) $f(\delta(q, a)) = \delta'(f(q), a)$ for all $q \in Q$ and $a \in \Sigma$;
- (2) $f(q_0) = q'_0$; and
- (3) $f(F) \subset F'$.

We say that M is a *binary subautomaton* of M' , if $Q \subset Q'$, and if the inclusion $i : Q \hookrightarrow Q'$ is a morphism of binary automata. This is equivalent to the assertion that $Q \subset Q'$, that δ is the restriction of δ' , that $q'_0 \in Q$, and that $F \subset F'$.

A *pointed binary graph* is a pair $P = (\Gamma, q_0)$, where $\Gamma = (Q, \delta)$ is a binary graph, and $q_0 \in Q$ is a distinguished node, called the *base node*. Let $P = (\Gamma, q_0)$ and $P' = (\Gamma', q'_0)$ be pointed binary graphs. A *morphism* from P to P' is a morphism of binary graphs $f : \Gamma \rightarrow \Gamma'$, such that $f(q_0) = q'_0$. We say that P is a *pointed binary subgraph* of P' if Γ is a binary subgraph of Γ' , and if the canonical monomorphism of binary graphs, $i : \Gamma \hookrightarrow \Gamma'$, is a morphism of pointed binary graphs. If $\Gamma = (Q, \delta)$ and $\Gamma' = (Q', \delta')$, this is equivalent to saying that $Q \subset Q'$, that δ is the restriction of δ' , and that $q'_0 \in Q$.

Given a binary automaton $M = (Q, \Sigma, \delta, q_0, F)$, we get a pointed binary graph $P_M = (\Gamma_M, q_0)$, where $\Gamma_M = (Q, \delta)$. The leaves of Γ_M are precisely the stationary states of M . In particular, if the automaton M is admissible, the leaves of Γ_M are the same as the final states of M . If $M = (Q, \Sigma, \delta, q_0, F)$, and $M' = (Q', \Sigma, \delta', q'_0, F')$ are two binary automata, and if $f : M \rightarrow M'$ is a morphism, then the underlying function $f : Q \rightarrow Q'$ defines a morphism of pointed binary graphs $P_f : P_M \rightarrow P_{M'}$. We thus obtain a functor p from the category $\mathcal{BA}$ of binary automata to the category $\mathcal{BG}_*$ of pointed binary graphs

$$\begin{aligned} p : \text{Ob}(\mathcal{BA}) &\rightarrow \text{Ob}(\mathcal{BG}_*), & p(M) &= P_M, & \text{and} \\ p : \text{Hom}(M, M') &\rightarrow \text{Hom}(P_M, P_{M'}), & p(f) &= P_f. \end{aligned}$$

We will see, in Section 11, that this morphism has quite an interesting structure.

Remark 7.1. The leaves of Γ_M are precisely the stationary states of M . In particular, if the automaton M is admissible, the leaves of Γ_M are the same as the final states of M . Moreover, the language $L(M) \subset \Sigma^*$ of an admissible automaton M is a monoidal ideal, i.e., if $w \in L(M)$ and $z \in \Sigma^*$, then $wz \in L(M)$.

In the theory of concurrent computation, one has the notion of a transition system (see [WN93, Chapter 2]). A *transition system* is a datum $T = (S, i, L, Tran)$, where

- (1) S is a set, whose elements are called *states*.
- (2) i is a distinguished element of S , called the *initial state*.
- (3) L is a set, whose elements are called *labels*.
- (4) $Tran$ is a subset of $S \times L \times S$, called the *transition relation*.

Let $T = (S, i, L, Tran)$ and $T' = (S', i', L', Tran')$ be transition systems. A *morphism* from T to T' is a pair (σ, λ) , where $\sigma : S \rightarrow S'$ is a function and $\lambda : L \dashrightarrow L'$ is a partial function (i.e., a function from a subset of L to L'), satisfying the following conditions:

- (1) $\sigma(i) = i'$.
- (2) If $(s_1, a, s_2) \in Tran$ and if $\lambda(a)$ is defined, then $(\sigma(s_1), \lambda(a), \sigma(s_2)) \in Tran'$.
- (3) If $(s_1, a, s_2) \in Tran$ and if $\lambda(a)$ is not defined, then $\sigma(s_1) = \sigma(s_2)$.

If $P = (\Gamma, q_0)$ is a pointed binary graph, with $\Gamma = (Q, \delta)$, we obtain a transition system $T_P = (Q, q_0, \Sigma, Gr(\delta))$, where $Gr(\delta) \subset Q \times \Sigma \times Q$ denotes the graph of δ , i.e.,

$$Gr(\delta) = \{(q_1, a, q_2) \in Q \times \Sigma \times Q \mid q_2 = \delta(q_1, a)\}.$$

Moreover, if $f : P \rightarrow P'$ is a morphism from a pointed binary graph $P = (\Gamma, q_0)$ to a pointed binary graph $P' = (\Gamma', q'_0)$, then the pair $(f, 1_\Sigma)$ is a morphism of the associated transition systems, $T_P \rightarrow T_{P'}$. This assignment provides us a functor $\mathcal{BG}_* \rightarrow \mathcal{TS}$, from the category of pointed binary graphs, to the category of transition systems. This functor is an isomorphism from $\mathcal{BG}_*$ to a subcategory of $\mathcal{TS}$. However, $\mathcal{BG}_*$ is not a full subcategory of $\mathcal{TS}$, as can be seen using simple examples of pointed binary graphs with two nodes.

8. FIBRED CATEGORIES

The original source for fibred categories is Grothendieck's article in SGA 1 [GR71, Exposé VI]. We also refer the reader to the notes of Streicher [Str99]. We recall below the basic notions about fibred categories. To aid the reader, and to make this article self-contained, we provide in this Section and the following two Sections, proofs of all propositions we need regarding fibred categories. The material is, perhaps, standard for specialists in certain areas of algebraic geometry and category theory. Such readers may proceed directly to Section 11.

Let $\mathcal{E}$ be a category. An $\mathcal{E}$ -category, or a *category over $\mathcal{E}$* , is a pair $(\mathcal{F}, p)$, where $\mathcal{F}$ is a category, and $p : \mathcal{F} \rightarrow \mathcal{E}$ is a functor. We usually drop the functor p from the notation, and say that $\mathcal{F}$ is an $\mathcal{E}$ -category. Sometimes, we also say that $p : \mathcal{F} \rightarrow \mathcal{E}$ is an $\mathcal{E}$ -category. An $\mathcal{E}$ -functor from an $\mathcal{E}$ -category $(\mathcal{F}, p)$ to an $\mathcal{E}$ -category $(\mathcal{G}, q)$ is a functor $f : \mathcal{F} \rightarrow \mathcal{G}$, such that $q \circ f = p$. In particular, a *section* of $(\mathcal{F}, p)$ is a functor $s : \mathcal{E} \rightarrow \mathcal{F}$ such that $p \circ s = 1_{\mathcal{E}}$.

Let $\mathcal{F}$ be an $\mathcal{E}$ -category, and let $S \in \text{Ob}(\mathcal{E})$. The *categorical fibre* of $\mathcal{F}$ at S is the subcategory $\mathcal{F}_S$ of $\mathcal{F}$ defined as follows:

- (1) $\text{Ob}(\mathcal{F}_S)$ is the collection of all $\xi \in \text{Ob}(\mathcal{F})$ such that $p(\xi) = S$.
- (2) If $\xi, \xi' \in \text{Ob}(\mathcal{F}_S)$, a morphism from ξ to ξ' in $\mathcal{F}_S$ is a morphism $u : \xi \rightarrow \xi'$ in $\mathcal{F}$, such that $p(u) = 1_S$. We will call such a morphism u an *S -morphism*, and we denote by $\text{Hom}_S(\xi, \xi')$, the set of all S -morphisms from ξ to ξ' .

We generalize the above notion of an $\mathcal{F}_S$ -morphism as follows. Suppose $f : T \rightarrow S$ is a morphism in $\mathcal{E}$, let $\eta \in \text{Ob}(\mathcal{F}_T)$, and let $\xi \in \text{Ob}(\mathcal{F}_S)$. An *f -morphism* from η to ξ is a morphism $u : \eta \rightarrow \xi$ in $\mathcal{F}$, such that $p(u) = f$. We denote the collection of all f -morphisms from η to ξ by $\text{Hom}_f(\eta, \xi)$.

Definition 8.1. (See [GR71, Exposé VI, Définition 5.1].) Let $(\mathcal{F}, p)$ be an $\mathcal{E}$ -category. Let $\alpha : \eta \rightarrow \xi$ be a morphism in $\mathcal{F}$, and let $S = p(\xi)$, $T = p(\eta)$ and $f = p(\alpha)$. We say that α is a *Cartesian*

morphism if for every object $\eta' \in \text{Ob}(\mathcal{F}_T)$, and for every f -morphism $u : \eta' \rightarrow \xi$, there exists a unique T -morphism $\bar{u} : \eta' \rightarrow \eta$ such that $u = \alpha \circ \bar{u}$. The pair (η, α) is called an *inverse image* of ξ by f .

The situation in the above definition can be described as in the following diagram:

$$\begin{array}{ccc}
 \eta' & \xrightarrow{u} & \xi \\
 \searrow \bar{u} & & \downarrow \alpha \\
 & \eta & \xrightarrow{\alpha} \xi
 \end{array}$$

$$\begin{array}{ccc}
 & & \downarrow p \\
 T & \xrightarrow{f} & S \\
 \searrow 1_T & & \downarrow f \\
 & T & \xrightarrow{f} S
 \end{array}$$

Thus, a morphism $\alpha : \eta \rightarrow \xi$ in $\mathcal{F}$ is Cartesian if and only if for every f -morphism $u : \eta' \rightarrow \xi$, the factorization $f = f \circ 1_T$ of $f = p(\alpha)$ in $\mathcal{E}$ lifts uniquely to a factorization $u = \alpha \circ \bar{u}$ of u in $\mathcal{F}$. In other words, $\alpha : \eta \rightarrow \xi$ is Cartesian if and only if for every $\eta' \in \text{Ob}(\mathcal{F}_T)$, the function

$$\begin{aligned}
 \alpha_* : \text{Hom}_T(\eta', \eta) &\rightarrow \text{Hom}_f(\eta', \xi) \\
 v &\mapsto \alpha \circ v
 \end{aligned}$$

is a bijection. We will use the notation α_* often, and record it for clarity.

Notation 8.2. Let $\mathcal{F}$ be a category, and let $\alpha : \eta \rightarrow \xi$ be a morphism in $\mathcal{F}$. Then, for every $\zeta \in \text{Ob}(\mathcal{F})$, we denote by α_* the function

$$\begin{aligned}
 \text{Hom}(\zeta, \eta) &\rightarrow \text{Hom}(\zeta, \xi), \\
 v &\mapsto \alpha \circ v.
 \end{aligned}$$

We denote the restriction of α_* to any subset F of $\text{Hom}(\zeta, \eta)$, also, by α_* .

Let $f : T \rightarrow S$ be a morphism in $\mathcal{E}$, and let $\xi \in \text{Ob}(\mathcal{F}_S)$. Suppose that (η, α) and (η', α') are two inverse images of ξ by f . Then, by definition, there exists a unique T -isomorphism $\bar{\alpha}' : \eta' \rightarrow \eta$ such that the diagram

$$\begin{array}{ccc}
 \eta' & \xrightarrow{\alpha'} & \xi \\
 \searrow \bar{\alpha}' & & \downarrow \alpha \\
 & \eta & \xrightarrow{\alpha} \xi
 \end{array}$$

commutes. Thus, an inverse image of ξ by f , if it exists, is unique up to a canonical T -isomorphism. Suppose that an inverse image of ξ by f exists, and that we have made a choice of such an inverse image. In such a situation, we often denote the chosen inverse image by $(f^*\xi, \alpha_f(\xi))$, or simply by $(f^*\xi, \alpha_f(\xi))$. Further, by abuse of language, we then call $(f^*\xi, \alpha_f(\xi))$ *the* inverse image of ξ by f . If an inverse image of ξ by f exists for all morphisms $f : T \rightarrow S$ in $\mathcal{E}$, and for all $\xi \in \text{Ob}(\mathcal{F}_S)$, then we say that *the inverse image functor by f in $\mathcal{F}$ exists*. If this is indeed the case, and if we have chosen, for all f and for all ξ , an inverse image of ξ by f , then, the assignment

$$\begin{aligned}
 f^* : \text{Ob}(\mathcal{F}_S) &\rightarrow \text{Ob}(\mathcal{F}_T), \quad \xi \mapsto f^*\xi \\
 f^* : \text{Hom}_S(\xi, \xi') &\rightarrow \text{Hom}_T(f^*\xi, f^*\xi'), \quad u \mapsto f^*u,
 \end{aligned}$$

is a functor from $\mathcal{F}_S$ to $\mathcal{F}_T$. Here, $f^*u : f^*\xi \rightarrow f^*\xi'$ is the unique T -morphism induced by the f -morphism $u \circ \alpha_f(\xi) : f^*\xi \rightarrow \xi'$, using the universal property of $f^*\xi'$.

Definition 8.3. (See [GR71, Exposé VI, Définition 6.1].) Let $\mathcal{E}$ be a category. We say that an $\mathcal{E}$ -category $(\mathcal{F}, p)$ is a *fibred category* over $\mathcal{E}$ if it satisfies the following two conditions:

- Fib_I** For every morphism $f : T \rightarrow S$ in $\mathcal{E}$, the inverse image functor by f in $\mathcal{F}$ exists.
- Fib_{II}** If $\alpha : \eta \rightarrow \xi$ and $\beta : \theta \rightarrow \eta$ are Cartesian morphisms in $\mathcal{F}$, then their composition, $\alpha \circ \beta : \theta \rightarrow \xi$, is also a Cartesian morphism.


We will see presently, in Proposition 9.3, that in fibred categories, Cartesian morphisms satisfy a certain stronger condition.

9. A CRITERION FOR FIBREDNESS

We will now present a criterion for a category $\mathcal{F}$ over $\mathcal{E}$ to be fibred. This criterion is sometimes taken to be a definition of fibred categories, as in [Str99, Definitions 2.1 and 2.2].

Definition 9.1. (See [WN93, Appendix B, page 163].) Let $(\mathcal{F}, p)$ be an $\mathcal{E}$ -category. Let $\alpha : \eta \rightarrow \xi$ be a morphism in $\mathcal{F}$, and let $S = p(\xi)$, $T = p(\eta)$ and $f = p(\alpha)$. We say that α is a *strongly Cartesian* morphism if for every morphism $g : U \rightarrow T$ in $\mathcal{E}$, for every object $\zeta \in \text{Ob}(\mathcal{F}_U)$, and for every $(f \circ g)$ -morphism $u : \zeta \rightarrow \xi$, there exists a unique g -morphism $\bar{u} : \zeta \rightarrow \eta$ such that $u = \alpha \circ \bar{u}$.

The situation in the above definition can be described as in the following diagram:



Thus, a morphism $\alpha : \eta \rightarrow \xi$ in $\mathcal{F}$ is strongly Cartesian if and only if for every morphism $g : U \rightarrow T$ in $\mathcal{E}$, for every $\zeta \in \text{Ob}(\mathcal{F}_U)$, and for every $(f \circ g)$ -morphism $u : \zeta \rightarrow \xi$, the factorization $f \circ g = f \circ g$ of $f \circ g$ in $\mathcal{E}$ lifts uniquely to a factorization $u = \alpha \circ \bar{u}$ of u in $\mathcal{F}$. In other words, $\alpha : \eta \rightarrow \xi$ is strongly Cartesian if and only if for every morphism $g : U \rightarrow T$ in $\mathcal{E}$, and for every $\zeta \in \text{Ob}(\mathcal{F}_U)$, the function

$$\begin{aligned} \alpha_* : \text{Hom}_g(\zeta, \eta) &\rightarrow \text{Hom}_{f \circ g}(\zeta, \xi) \\ v &\mapsto \alpha \circ v \end{aligned}$$

is a bijection.

Strongly Cartesian morphisms behave well under composition.

Proposition 9.2. (See [WN93, Appendix B, page 163].) Let $\mathcal{E}$ be a category, and $(\mathcal{F}, p)$ an $\mathcal{E}$ -category. Suppose $\alpha : \eta \rightarrow \xi$ and $\beta : \zeta \rightarrow \eta$ are strongly Cartesian morphisms in $\mathcal{F}$. Then, their composition $\alpha \circ \beta : \zeta \rightarrow \xi$ is also strongly Cartesian.

Proof. Let $f : T \rightarrow S$ and $g : U \rightarrow T$ denote $p(\alpha)$ and $p(\beta)$, respectively. Let $h : V \rightarrow U$ be a morphism in $\mathcal{E}$, and let $\theta \in \text{Ob}(\mathcal{F}_V)$. Then, the diagram

$$\begin{array}{ccc} \text{Hom}_h(\theta, \zeta) & \xrightarrow{\beta_*} & \text{Hom}_{g \circ h}(\theta, \eta) \\ & \searrow (\alpha \circ \beta)_* & \downarrow \alpha_* \\ & & \text{Hom}_{f \circ g \circ h}(\theta, \xi) \end{array}$$

commutes. Since α and β are strongly Cartesian morphisms, the functions α_* and β_* are bijections. Therefore, by the commutativity of the above diagram, the function $(\alpha \circ \beta)_*$ is also a bijection and, hence, $\alpha \circ \beta$ is strongly Cartesian. $\square$

Taking $g = 1_T$ in Definition 9.1, we see that every strongly Cartesian morphism is Cartesian. The converse also is true in fibred categories.

Proposition 9.3. (See [GR71, Exposé VI, Proposition 6.11].) *Let $\mathcal{E}$ be a category, and $(\mathcal{F}, p)$ an $\mathcal{E}$ -category. Then, $\mathcal{F}$ is a fibred category over $\mathcal{E}$ if and only if it satisfies the condition **Fib_I** of Definition 8.3, and the following condition:*

Fib'_{II} Every Cartesian morphism in $\mathcal{F}$ is strongly Cartesian.

Proof. Suppose that $\mathcal{F}$ satisfies condition **Fib_I** of Definition 8.3. Then, we need to show that $\mathcal{F}$ satisfies **Fib_{II}** if and only if it satisfies **Fib'_{II}**.

So, let us suppose, first, that $\mathcal{F}$ satisfies **Fib_{II}**. We will prove that every Cartesian morphism $\alpha : \eta \rightarrow \xi$ in $\mathcal{F}$ is strongly Cartesian. Let $f : T \rightarrow S$, $g : U \rightarrow T$ and $\zeta \in \text{Ob}(\mathcal{F}_U)$ be as in Definition 9.1. Let $\beta : \zeta' \rightarrow \eta$ be the inverse image of η by g . Then, the diagram

$$\begin{array}{ccc} \text{Hom}_U(\zeta, \zeta') & \xrightarrow{\beta_*} & \text{Hom}_g(\zeta, \eta) \\ & \searrow (\alpha \circ \beta)_* & \downarrow \alpha_* \\ & & \text{Hom}_{f \circ g}(\zeta, \xi) \end{array}$$

commutes. Since (ζ', β) is the inverse image of an object, the morphism β is Cartesian and, hence, the function β_* is a bijection. On the other hand, by **Fib_{II}**, the composition $\alpha \circ \beta : \zeta' \rightarrow \xi$ is a Cartesian morphism. Therefore, the function $(\alpha \circ \beta)_*$ is also a bijection. Thus, by the commutativity of the above diagram, the function α_* is a bijection, as well. In other words, α is a strongly Cartesian morphism.

Conversely, if $\mathcal{F}$ satisfies **Fib'_{II}**, then every composition of Cartesian morphisms is a product of strongly Cartesian morphisms, which, by Proposition 9.2, is strongly Cartesian, hence Cartesian. Therefore, $\mathcal{F}$ satisfies the condition **Fib_{II}**. $\square$

10. SPLIT CATEGORIES

We will now discuss a special class of fibred categories, which is relevant to our description of binary graphs. We refer the reader to [GR71, Exposé VI, Sections 7, 8 and 9] for a discussion, in full generality, of the topics mentioned in this Section.

Definition 10.1. (See [GR71, Exposé VI, Définition 7.1].) Let $\mathcal{F}$ be an $\mathcal{E}$ -category. A *cleavage* of $\mathcal{F}$ over $\mathcal{E}$ is an attachment, to each morphism $f : T \rightarrow S$ in $\mathcal{E}$, of an inverse image functor by f in $\mathcal{F}$, say, f^* . A *cleaved category* over $\mathcal{E}$ is an $\mathcal{E}$ -category $\mathcal{F}$, together with a cleavage of $\mathcal{F}$.

Definition 10.2. (See [GR71, Exposé VI, Section 9], and [Str99, Definition 4.1].) Let $\mathcal{F}$ be an $\mathcal{E}$ -category. We say that a cleavage of $\mathcal{F}$ over $\mathcal{E}$ is a *splitting* if it satisfies the following conditions:

- (1) For every $S \in \text{Ob}(\mathcal{E})$, the inverse image functor $(1_S)^* : \mathcal{F}_S \rightarrow \mathcal{F}_S$ equals the identity functor $1_{\mathcal{F}_S}$, and for every object $\xi \in \text{Ob}(\mathcal{F}_S)$, we have $\alpha_{1_S}(\xi) = 1_\xi$, where $\alpha_{1_S}(\xi) : (1_S)^*\xi \rightarrow \xi$ is the canonical morphism.
- (2) For every pair of composable morphisms, say, $f : T \rightarrow S$ and $g : U \rightarrow T$ in $\mathcal{E}$, we have

$$(f \circ g)^* = g^* \circ f^* : \mathcal{F}_S \rightarrow \mathcal{F}_U,$$

and for all $\xi \in \text{Ob}(\mathcal{F}_S)$, we have

$$\alpha_{f \circ g}(\xi) = \alpha_f(\xi) \circ \alpha_g(f^*\xi) : (f \circ g)^*\xi = g^*f^*\xi \rightarrow \xi,$$

where $\alpha_f(\xi) : f^*\xi \rightarrow \xi$ is, as usual, the canonical morphism.

A *split category* over $\mathcal{E}$ is an $\mathcal{E}$ -category $\mathcal{F}$, together with a splitting of $\mathcal{F}$. If $\mathcal{F}$ and $\mathcal{G}$ are split categories over $\mathcal{E}$, then, a *morphism of split categories*, from $\mathcal{F}$ to $\mathcal{G}$, is an $\mathcal{E}$ -functor $F : \mathcal{F} \rightarrow \mathcal{G}$, such that for every morphism $f : T \rightarrow S$ in $\mathcal{E}$, and for every $\xi \in \text{Ob}(\mathcal{F}_S)$, we have $F(f^*\xi) = f^*F(\xi)$ and $F(\alpha_f(\xi)) = \alpha_f(F(\xi))$. We thus obtain a category $\mathbf{Split}_{\mathcal{E}}$ of split categories over $\mathcal{E}$.

Notation 10.3. Let $\mathbf{Cat}$ denote the “category of categories”, i.e., the category whose objects are categories, and whose morphisms are functors between categories. For any pair of categories $\mathcal{A}$ and $\mathcal{B}$, let $\mathbf{Hom}(\mathcal{A}, \mathcal{B})$ denote the category whose objects are functors from $\mathcal{A}$ to $\mathcal{B}$, and whose morphisms are natural transformations of functors. In particular, $\mathbf{Hom}(\mathcal{A}^\circ, \mathcal{B})$, where $\mathcal{A}^\circ$ is the opposite category of $\mathcal{A}$, is the category of contravariant functors from $\mathcal{A}$ to $\mathcal{B}$.

Remark 10.4. Given a split category $\mathcal{F}$ over $\mathcal{E}$, we obtain a functor $\phi(\mathcal{F}) : \mathcal{E}^\circ \rightarrow \mathbf{Cat}$, that is defined as follows:

$$\begin{aligned} \phi(\mathcal{F}) : \text{Ob}(\mathcal{E}^\circ) &\rightarrow \text{Ob}(\mathbf{Cat}), & S &\mapsto \mathcal{F}_S, & \text{and} \\ \phi(\mathcal{F}) : \text{Hom}(T, S) &\rightarrow \text{Hom}(\mathcal{F}_S, \mathcal{F}_T), & f &\mapsto f^*. \end{aligned}$$

If $F : \mathcal{F} \rightarrow \mathcal{G}$ is a morphism in $\mathbf{Split}_{\mathcal{E}}$, we obtain a natural transformation $\phi(F) : \phi(\mathcal{F}) \rightarrow \phi(\mathcal{G})$ as follows: given any object $S \in \text{Ob}(\mathcal{E})$, we define

$$\phi(F)_S : \phi(\mathcal{F})(S) = \mathcal{F}_S \rightarrow \phi(\mathcal{G})(S) = \mathcal{G}_S, \quad \phi(F)_S = F|_{\mathcal{F}_S}.$$

We thus obtain a functor

$$\phi : \mathbf{Split}_{\mathcal{E}} \rightarrow \mathbf{Hom}(\mathcal{E}^\circ, \mathbf{Cat}).$$

This functor, ϕ , is an equivalence of categories (see [GR71, Exposé VI, Section 9]).

Proposition 10.5. *Every split category $\mathcal{F}$, over a category $\mathcal{E}$, is a fibred category.*

Proof. Since $\mathcal{F}$ admits a cleavage, it satisfies the condition $\mathbf{Fib}_I$ of Definition 8.3. We will show that $\mathcal{F}$ satisfies the condition $\mathbf{Fib}'_{II}$ of Proposition 9.3, i.e., that every Cartesian morphism in $\mathcal{F}$ is strongly Cartesian. Let $\alpha : \eta \rightarrow \xi$ be a Cartesian morphism in $\mathcal{F}$, and let $f : T \rightarrow S$ denote the morphism $p(f)$ in $\mathcal{E}$, where $p : \mathcal{F} \rightarrow \mathcal{E}$ is the canonical functor. Let $g : U \rightarrow T$ be a morphism in $\mathcal{E}$, and let $\zeta \in \text{Ob}(\mathcal{F}_U)$. We, then, have to show that the function

$$\begin{aligned} \alpha_* : \text{Hom}_g(\zeta, \eta) &\rightarrow \text{Hom}_{f \circ g}(\zeta, \xi), \\ v &\mapsto \alpha \circ v, \end{aligned}$$

is a bijection. Since α is Cartesian, the pair (η, α) is an inverse image of ξ by f . On the other hand, the given splitting of $\mathcal{F}$ provides another inverse image $(f^*\xi, \alpha_f(\xi))$ of ξ by f . Therefore, by the definition

of Cartesian morphisms, there exists a unique T -isomorphism $\beta : f^*\xi \rightarrow \eta$ such that $\alpha_f(\xi) = \alpha \circ \beta$. We now have the following commutative diagram:

$$\begin{array}{ccc}
 \mathrm{Hom}_g(\zeta, \eta) & \xrightarrow{\alpha_*} & \mathrm{Hom}_{f \circ g}(\zeta, \xi) \\
 \uparrow \beta_* & & \uparrow \alpha_{f \circ g}(\xi)_* \\
 \mathrm{Hom}_g(\zeta, f^*\xi) & \xleftarrow{\alpha_g(f^*\xi)_*} & \mathrm{Hom}_U(\zeta, (f \circ g)^*\xi) \\
 & \nwarrow \alpha_g(f^*\xi)_* & \nearrow \\
 & \mathrm{Hom}_U(\zeta, g^*f^*\xi) &
 \end{array}$$

The triangle in the bottom of the diagram is defined because $g^*f^*\xi = (f \circ g)^*\xi$, and the commutativity of the rectangle follows from the equations

$$\alpha \circ \beta \circ \alpha_g(f^*\xi) = \alpha_f(\xi) \circ \alpha_g(f^*\xi) = \alpha_{f \circ g}(\xi).$$

The function β_* is a bijection because β is an isomorphism, while the bottom and right sides of the rectangle are bijections because the morphisms $\alpha_g(f^*\xi)$ and $\alpha_{f \circ g}(\xi)$, respectively, are Cartesian. Therefore, the top edge of the rectangle, too, is a bijection. This proves that $\alpha : \eta \rightarrow \xi$ is strongly Cartesian. $\square$

Definition 10.6. (See [GR71, Exposé VI, Section 9].) We say that a category $\mathcal{A}$ is *rigid* if, for every object $\xi \in \mathrm{Ob}(\mathcal{A})$, the identity morphism 1_ξ is the only automorphism of ξ . The category $\mathcal{A}$ is said to be *reduced*, if whenever two objects in $\mathcal{A}$ are isomorphic, they are, in fact, equal.

Proposition 10.7. Let $\mathcal{E}$ be a category, and let $\mathcal{F}$ be an $\mathcal{E}$ -category. Suppose that for every object $S \in \mathrm{Ob}(\mathcal{E})$, the categorical fibre $\mathcal{F}_S$ is a rigid and reduced category. Then, $\mathcal{F}$ is a fibred category if and only if it is split. If this is, indeed, the case, then there exists a unique cleavage of $\mathcal{F}$ over $\mathcal{E}$, and that cleavage is a splitting.

Proof. By Proposition 10.5, if $\mathcal{F}$ is split over $\mathcal{E}$, then it is a fibred category. Conversely, suppose that $\mathcal{F}$ is a fibred category over $\mathcal{E}$. We will, then, show that $\mathcal{F}$ has a unique cleavage, and that this cleavage is a splitting of $\mathcal{F}$ over $\mathcal{E}$.

Since $\mathcal{F}$ is a fibred category over $\mathcal{E}$, using the Axiom of Choice, as in [Str99, Section 4], we see that $\mathcal{F}$ admits a cleavage. Suppose that $\mathcal{F}$ admits two cleavages $*$ and $\dagger$. For any morphism $f : T \rightarrow S$ in $\mathcal{E}$, and for any object $\xi \in \mathrm{Ob}(\mathcal{F}_S)$, denote the inverse images of ξ by f , with respect to the cleavages $*$ and $\dagger$, by $(f^*\xi, \alpha_f(\xi))$ and $(f^\dagger\xi, \beta_f(\xi))$, respectively. Then, there exists a unique T -isomorphism $\theta_f(\xi) : f^*\xi \rightarrow f^\dagger\xi$, such that $\alpha_f(\xi) = \beta_f(\xi) \circ \theta_f(\xi)$. Since $\mathcal{F}_T$ is a reduced category, the T -isomorphic objects $f^*\xi$ and $f^\dagger\xi$ are equal. Now, because $\mathcal{F}_T$ is a rigid category, the T -automorphism $\theta_f(\xi)$ of $f^*\xi$ equals the identity morphism. Therefore, $\alpha_f(\xi) = \beta_f(\xi)$. In other words, for all f and ξ , the inverse images $(f^*\xi, \alpha_f(\xi))$ and $(f^\dagger\xi, \beta_f(\xi))$ are equal. Thus, the two cleavages $*$ and $\dagger$ are equal and, so, $\mathcal{F}$ has a unique cleavage over $\mathcal{E}$.

We will show, next, that the unique cleavage of $\mathcal{F}$ is a splitting. Let f and ξ be as above, and let $g : U \rightarrow T$ be another morphism in $\mathcal{E}$. Then, by the condition **Fib_{II}** of Definition 8.3, the composition $\alpha_f(\xi) \circ \alpha_g(f^*\xi) : g^*f^*\xi \rightarrow \xi$ is a Cartesian morphism over $f \circ g : U \rightarrow S$. Therefore, by the universal property of inverse images, there exists a unique U -isomorphism $c_{f,g}(\xi) : g^*f^*\xi \rightarrow (f \circ g)^*\xi$, such

that the following diagram commutes:

$$\begin{array}{ccc}
 g^* f^* \xi & \xrightarrow{\alpha_g(f^* \xi)} & f^* \xi \\
 \downarrow c_{f,g}(\xi) & & \downarrow \alpha_f(\xi) \\
 (f \circ g)^* \xi & \xrightarrow{\alpha_{f \circ g}(\xi)} & \xi
 \end{array}$$

Because $\mathcal{F}_U$ is a rigid and reduced category, we know that the U -isomorphic objects $g^* f^* \xi$ and $(f \circ g)^* \xi$ are equal, and that the U -automorphism $c_{f,g}(\xi)$ of $(f \circ g)^* \xi$ is the identity morphism. Therefore, we get $\alpha_{f \circ g}(\xi) = \alpha_f(\xi) \circ \alpha_g(f^* \xi)$. In a similar manner, since $(\xi, 1_\xi)$ and $((1_S)^* \xi, \alpha_{1_S}(\xi))$ are both inverse images of ξ by 1_S , we see that $((1_S)^* \xi, \alpha_{1_S}(\xi)) = (\xi, 1_\xi)$. We conclude that the unique cleavage of $\mathcal{F}$ is a splitting over $\mathcal{E}$. $\square$

11. FIBRED CATEGORY OF BINARY AUTOMATA

Recall that in Section 7, we have defined a functor $p : \mathcal{BA} \rightarrow \mathcal{BG}_*$, from the category $\mathcal{BA}$ of binary automata to the category $\mathcal{BG}_*$ of pointed binary graphs. In the terminology of Section 8, $(\mathcal{BA}, p)$ is a $\mathcal{BG}_*$ -category. We will now show that it is, in fact, a fibred category over $\mathcal{BG}_*$. But, first, let us note that we have a canonical functor in the other direction, too.

Given a pointed binary graph $P = (\Gamma, q_0)$, where $\Gamma = (Q, \delta)$ we get an admissible binary automaton $M_P = (Q, \Sigma, \delta, q_0, F)$, where F is the set of leaves in Q . If $P = (\Gamma, q_0)$, and $P' = (\Gamma', q'_0)$ are two pointed binary graphs, and if $f : P \rightarrow P'$ is a morphism, then the underlying function $f : Q \rightarrow Q'$, being Σ^* -equivariant, takes the leaves of Γ to leaves of Γ' and, so, defines a morphism of binary automata $M_f : M_P \rightarrow M_{P'}$. We thus obtain a functor s from the category $\mathcal{BG}_*$ to the category $\mathcal{BA}$,

$$\begin{aligned}
 s : \text{Ob}(\mathcal{BG}_*) &\rightarrow \text{Ob}(\mathcal{BA}), & s(P) &= M_P, & \text{and} \\
 s : \text{Hom}(P, P') &\rightarrow \text{Hom}(M_P, M_{P'}), & s(f) &= M_f.
 \end{aligned}$$

Proposition 11.1. *The above functor $s : \mathcal{BG}_* \rightarrow \mathcal{BA}$ is a section of the $\mathcal{BG}_*$ -category $(\mathcal{BA}, p)$. This section is an isomorphism of categories from $\mathcal{BG}_*$, to the full subcategory of $\mathcal{BA}$ consisting of admissible automata.*

Proof. It is clear that $p \circ s = 1_{\mathcal{BG}_*}$. From the definition of admissibility, it follows that for every pointed binary graph P , the binary automaton M_P is admissible. Let $M = (Q, \Sigma, \delta, q_0, F)$ be an admissible automaton, and let $P_M = (\Gamma_M, q_0)$. Then, it follows, from Remark 7.1, that the set of leaves of Γ_M equals the set, F , of final states of M . Thus $s(P_M) = M$, i.e., $s \circ p = 1$ on $\text{Ob}(\mathcal{ABA})$, where $\mathcal{ABA}$ is the full subcategory of $\mathcal{BA}$, consisting of admissible binary automata. Moreover, if $f : M \rightarrow M'$ is a morphism in $\mathcal{ABA}$, then $s \circ p(f) = s(P_f) = M_{P_f} = f$. Thus, $s \circ p = 1$ on $\text{Hom}(M, M')$, and hence $s \circ p = 1_{\mathcal{ABA}}$. $\square$

We, thus, obtain an identification between pointed binary graphs and admissible binary automata.

Remark 11.2. It follows, from Proposition 11.1, that $s : \text{Ob}(\mathcal{BG}_*) \rightarrow \text{Ob}(\mathcal{BA})$ is an injective function. On the other hand, $p : \text{Ob}(\mathcal{BA}) \rightarrow \text{Ob}(\mathcal{BG}_*)$ is a many-to-one function. Indeed if $P = (\Gamma, q_0) \in \text{Ob}(\mathcal{BG}_*)$, and if $\Gamma = (Q, \delta)$, then for every subset $F \subset Q$, we obtain an object $M = (Q, \Sigma, \delta, q_0, F)$ such that $p(M) = P$. Indeed, the categorical fibre, $\mathcal{BA}_P$, of $\mathcal{BA}$ over P can be described as follows:

- (1) $\text{Ob}(\mathcal{BA}_P) = 2^Q$, the power set of Q .
- (2) If F_1 and F_2 are subsets of Q , then $\text{Hom}_P(F_1, F_2)$ is the empty set if F_1 is not a subset of F_2 , and is the singleton $\{\iota_{F_1, F_2}\}$, if $F_1 \subset F_2$, in which case ι_{F_1, F_2} is the inclusion map $F_1 \hookrightarrow F_2$.

We call this category the *posetal category defined by the power set of Q* , and denote it by $\mathcal{P}_Q$.

Definition 11.3. Let $P = (\Gamma, q_0)$, and $P' = (\Gamma', q'_0)$ be two pointed binary graphs, where $\Gamma = (Q, \delta)$ and $\Gamma' = (Q', \delta')$, and let $f : P \rightarrow P'$ be a morphism. Let $M' \in \text{Ob}(\mathcal{BA}_{P'})$ be a binary automaton over P' , say, $M' = (Q', \Sigma, \delta', q'_0, F')$. We, then, define a binary automaton $f^*M' \in \text{Ob}(\mathcal{BA}_P)$ by $f^*M' = (Q, \Sigma, \delta, q_0, f^{-1}(F'))$. Further, we define a morphism of binary automata, $\alpha_f(M') : f^*M' \rightarrow M'$ by setting $\alpha_f(M') = f : Q \rightarrow Q'$. We will show below (see the proof of Theorem 11.4), that the pair $(f^*M', \alpha_f(M'))$ is an inverse image of M' by f . We thus obtain a cleavage of $\mathcal{BA}$ over $\mathcal{BG}_*$. We call it the *canonical cleavage* of $\mathcal{BA}$ over $\mathcal{BG}_*$.

Theorem 11.4. *The $\mathcal{BG}_*$ -category $(\mathcal{BA}, p)$ is a fibred category, in fact, a split category. The canonical cleavage of $\mathcal{BA}$ (see Definition 11.3) is its only cleavage, and this cleavage is a splitting.*

Proof. We will, first, prove that the canonical cleavage is, indeed, a cleavage of $\mathcal{BA}$ over $\mathcal{BG}_*$. Let $P = (\Gamma, q_0)$, and $P' = (\Gamma', q'_0)$ be two pointed binary graphs, where $\Gamma = (Q, \delta)$ and $\Gamma' = (Q', \delta')$, and let $f : P \rightarrow P'$ be a morphism. Let $M' \in \text{Ob}(\mathcal{BA}_{P'})$ be a binary automaton over P' , say, $M' = (Q', \Sigma, \delta', q'_0, F')$. Let $(f^*M', \alpha_f(M'))$ be as in Definition 11.3. Let $M'' \in \text{Ob}(\mathcal{BA}_P)$ be a binary automaton over P , say, $M'' = (Q, \Sigma, \delta, q_0, F'')$, and let $u : M'' \rightarrow M'$ be an f -morphism of binary automata. We need to show that there exists a unique P -morphism $\bar{u} : M'' \rightarrow f^*M'$ such that the diagram

$$\begin{array}{ccc} M'' & & \\ & \searrow u & \\ & f^*M' & \xrightarrow{\alpha_f(M')} M' \\ & \nearrow \bar{u} & \end{array}$$

commutes. Since u is an f -morphism, by the definition of the functor $p : \mathcal{BA} \rightarrow \mathcal{BG}_*$, we see that the underlying function $u : Q \rightarrow Q'$ of u , equals f . Further, since u is a morphism of binary automata, we have $f(F'') = u(F'') \subset F'$, i.e., we have $F'' \subset f^{-1}(F')$. Now, define $\bar{u} : M'' \rightarrow f^*M'$, by $\bar{u} = 1_Q : Q \rightarrow Q$. Then, the fact that F'' is a subset of $f^{-1}(F')$ implies that $\bar{u}$ is a morphism of binary automata. It is, obviously, a P -morphism and, moreover,

$$\alpha_f(M') \circ \bar{u} = f \circ 1_Q = f = u.$$

This proves the “existence” part of Cartesianness. To prove the “uniqueness” part, suppose that v and w are P -morphisms from M'' to f^*M' , such that $\alpha_f(M') \circ v = \alpha_f(M') \circ w$. Since $p(v) = 1_P$, the underlying function $v : Q \rightarrow Q$ equals 1_Q . Similarly, w equals 1_Q , hence $v = w$. We have, thus, proved that the canonical cleavage is, in fact, a cleavage of $\mathcal{BA}$ over $\mathcal{BG}_*$.

We will, next, show that the canonical cleavage of $\mathcal{BA}$ is a splitting. If $f = 1_{P'}$, then $f^{-1}(F') = F'$, hence $(1_{P'})^*M' = M'$, and $\alpha_{1_{P'}}(M') = 1_{M'}$. Next, suppose that f and M' are as above, and let $g : P'' \rightarrow P$ be another morphism in $\mathcal{BG}_*$. Then, $(f \circ g)^{-1}(F') = g^{-1}(f^{-1}(F'))$, hence $(f \circ g)^*M' = g^*f^*M'$. Further, since the underlying function of a morphism of the type $\alpha_f(M')$, as we saw above, equals f , we have

$$\alpha_{f \circ g}(M') = f \circ g = \alpha_f(M') \circ \alpha_g(f^*M').$$

Therefore, the canonical cleavage of $\mathcal{BA}$ is a splitting over $\mathcal{BG}_*$. Thus, $(\mathcal{BA}, p)$ is a split category and, so, by Proposition 10.5, it is a fibred category over $\mathcal{BG}_*$.

Finally, we have observed (see Remark 11.2), that for every object $P \in \text{Ob}(\mathcal{BG}_*)$, the categorical fibre $\mathcal{BA}_P$ is the posetal category $\mathcal{P}_Q$ defined by the power set of Q , where Q is the set of nodes of P . Thus, every categorical fibre of $\mathcal{BA}$ is a rigid and reduced category. Therefore, by Proposition 10.7, the fibred category $\mathcal{BA}$ has a unique cleavage. $\square$

Remark 11.5. It follows from Remark 10.4 that there is a canonical equivalence ϕ from the category $\text{Split}_{\mathcal{BG}_*}$ to the category $\text{Hom}(\mathcal{BG}_*^\circ, \text{Cat})$. From the definition of ϕ , we see that the functor $\phi(\mathcal{BA}) : \mathcal{BG}_*^\circ \rightarrow \text{Cat}$ is isomorphic to the following functor:

$$\begin{aligned} \text{Ob}(\mathcal{BG}_*^\circ) &\rightarrow \text{Cat}, & P &\mapsto \mathcal{P}_Q, & \text{and} \\ \text{Hom}(P, P') &\rightarrow \text{Hom}(\mathcal{P}_{Q'}, \mathcal{P}_Q), & f &\mapsto f^*. \end{aligned}$$

We use here the notation Q (respectively, Q') for the set of nodes of the pointed binary graph P (respectively, P'); if Q is a set, $\mathcal{P}_Q$ denotes the posetal category defined by the power set of Q (see Remark 11.2); and, for any function $f : Q \rightarrow Q'$, the symbol f^* denotes the obvious pull-back functor from $\mathcal{P}_{Q'}$ to $\mathcal{P}_Q$.

12. CONCLUSION

In this article, we have developed a set-theoretic formalism for binary graphs. We have expressed various notions regarding graphs in terms of the inheritance order on the set of nodes, and in terms of the action of bit strings on nodes. We have exhibited pointed binary graphs as a subcategory of transition systems. Together with the set-theoretic notion of binary graphs, another interesting result, for us, in this paper is the fact that binary automata form a fibred category over pointed binary graphs. We would like to have a better understanding of this fibred category.

We suggest that it would be interesting to formulate various algorithms for trees, using the formalism developed here. Obviously, this formalism, alone, would not affect the efficiency of such algorithms. However, we feel that the abstract set-theoretic descriptions provided here would help in specifying the algorithms in a clear and precise manner.

ACKNOWLEDGEMENTS

I thank M. Vidyasagar for providing me an opportunity to interact with computational biologists, and Sharmila Mande and Ganesh Gayatri for introducing me to the algorithms that motivated this work. I am grateful to K. Viswanath for his comments on an earlier version of this article. I thank I.B.S. Passi for asking me to lecture in his seminar on geometric group theory, at the Harish-Chandra Research Institute, Allahabad; the talks I gave there, on automatic groups, motivated me to study finite automata.

REFERENCES

- [AGM95] S. Abramsky, D.M. Gabbay, and T.S.E Maibaum (eds.), *Handbook of logic in computer science, Vol. 4, Semantic modelling*, Oxford University Press, Oxford, UK, 1995, ISBN 0-19-853780-8.
- [AHU74] Alfred V. Aho, John E. Hopcroft, and Jeffrey D. Ullman, *The design and analysis of computer algorithms*, Addison-Wesley, Boston 1974, ISBN 981-4053-19-8.
- [GR71] Alexandre Grothendieck and Michele Raynaud, *Revêtements étales et groupe fondamental (SGA 1)*, *Un Séminaire dirigé par A. Grothendieck, Augmenté de deux exposés par M. Raynaud, Séminaire de Géométrie Algébrique du Bois-Marie, 1960-61*, Lecture Notes in Mathematics, no. 224, Springer-Verlag, Berlin, Heidelberg and New York, 1971, arXiv:math.AG/0206203, ISBN 3-540-05614-9.
- [HU79] John E. Hopcroft and Jeffrey D. Ullman, *Introduction to automata theory, languages, and computation*, first ed., Addison-Wesley, Boston, 1979, ISBN 0-201-02988-X.
- [Knu97] Donald E. Knuth, *The art of computer programming, Vol. 1, Fundamental algorithms*, third ed., Addison-Wesley, Boston, 1997, ISBN 81-7808-111-3.
- [Str99] Thomas Streicher, *Fibred categories à la J. Bénabou*, Notes of a course given at a spring school in Munich, URL <http://www.mathematik.tu-darmstadt.de/~streicher/>, 1999.
- [WN93] Glynn Winskel and Mogens Nielsen, *Models for concurrency*, Tech. Report PB-463, Department of Computer Science, University of Aarhus, Aarhus, Denmark, November 1993, URL <http://www.daimi.au.dk/PB/463/>, Preprint version of [WN95].
- [WN95] ———, *Models for concurrency*, in Abramsky et al. [AGM95], pp. 1–148.

ADVANCED TECHNOLOGY CENTRE, TATA CONSULTANCY SERVICES, 5-9-62 (6TH FLOOR), KHAN LATEEF KHAN ESTATE, FATEH MAIDAN ROAD, HYDERABAD 500 001, INDIA

E-mail address: raghu@atc.tcs.co.in